

Apparent Singularities of D-finite Systems

Yi Zhang

Johann Radon Institute for Computational and Applied Mathematics (RICAM)
Austrian Academy of Sciences, Austria

Joint work with Shaoshi Chen, Manuel Kauers and Ziming Li



Der Wissenschaftsfonds.

Singularities (univariate case)

Let $\partial = \frac{d}{dx}$.

Consider

$$L = p_r \partial^r + p_{r-1} \partial^{r-1} + \cdots + p_0 \in \mathbb{C}[x][\partial],$$

where $p_i \in \mathbb{C}[x]$ with $p_r \neq 0$ and $\gcd(p_r, p_{r-1}, \dots, p_0) = 1$.

Call r the **order** of L , denoted by $\text{ord}(L)$.

Singularities (univariate case)

Let $\partial = \frac{d}{dx}$.

Consider

$$L = p_r \partial^r + p_{r-1} \partial^{r-1} + \cdots + p_0 \in \mathbb{C}[x][\partial],$$

where $p_i \in \mathbb{C}[x]$ with $p_r \neq 0$ and $\gcd(p_r, p_{r-1}, \dots, p_0) = 1$.

Call r the **order** of L , denoted by $\text{ord}(L)$.

Definition. $c \in \mathbb{C}$ is an **ordinary point** of L if $p_r(c) \neq 0$.

Otherwise, c is a **singularity** of L .

Formal power series (univariate case)

Definition. Let $f \in \mathbb{C}[[x]]$ be of the form

$$f = c_m x^m + c_{m+1} x^{m+1} + \dots,$$

where $c_m \neq 0$. Call m the **initial exponent** of f .

Formal power series (univariate case)

Definition. Let $f \in \mathbb{C}[[x]]$ be of the form

$$f = c_m x^m + c_{m+1} x^{m+1} + \dots,$$

where $c_m \neq 0$. Call m the **initial exponent** of f .

Theorem (Fuchs, 1866). Let $L \in \mathbb{C}[x][\partial] \setminus \{0\}$. Then

the origin is an ordinary point of L



L has $\text{ord}(L)$ sols in $\mathbb{C}[[x]]$ with initial exponents $0, 1, \dots, \text{ord}(L) - 1$.

Apparent singularities

Assume the origin is a singularity of L .

Definition. The origin is **apparent** if L has $\text{ord}(L)$ \mathbb{C} -linearly independent sols in $\mathbb{C}[[x]]$.

Apparent singularities

Assume the origin is a singularity of L .

Definition. The origin is **apparent** if L has $\text{ord}(L)$ \mathbb{C} -linearly independent sols in $\mathbb{C}[[x]]$.

Example. x^5 is a sol of $xf'(x) - 5f(x) = 0$.

Motivation

Assume the origin is an apparent singularity of L .

Goal. Find $M \in \mathbb{C}[x][\partial] \setminus \{0\}$ s.t.

- ▶ $\text{sol}(L) \subset \text{sol}(M)$;
- ▶ the origin is an ordinary point of M .

Motivation

Assume the origin is an apparent singularity of L .

Goal. Find $M \in \mathbb{C}[x][\partial] \setminus \{0\}$ s.t.

- ▶ $\text{sol}(L) \subset \text{sol}(M)$;
- ▶ the origin is an ordinary point of M .

Remark. If so, then $\text{sol}(L)$ is spanned by formal power series.

Apparent singularities

L has sols of the form:

$$x^{\alpha_1} + \dots,$$

$$x^{\alpha_2} + \dots,$$

$$\vdots$$

$$x^{\alpha_r} + \dots.$$

where $\alpha_1 < \alpha_2 < \dots < \alpha_r \in \mathbb{N}$, $r = \text{ord}(L)$.

Remark. Some exponents are missing!

Apparent singularities

L has sols of the form:

$$\begin{aligned}x^{e_1} &+ \cdots, & e_1 &= 0, \dots, \alpha_1 - 1, \\x^{\alpha_1} &+ \cdots, \\x^{e_2} &+ \cdots, & e_2 &= \alpha_1 + 1, \dots, \alpha_2 - 1, \\x^{\alpha_2} &+ \cdots, \\&\vdots \\x^{e_r} &+ \cdots, & e_r &= \alpha_{r-1} + 1, \dots, \alpha_r - 1, \\x^{\alpha_r} &+ \cdots.\end{aligned}$$

where $\alpha_1 < \alpha_2 < \cdots < \alpha_r \in \mathbb{N}$, $r = \text{ord}(L)$.

Remark. Some exponents are missing!

Desingularization

Given $L \in \mathbb{C}[x][\partial]$, the origin being apparent, find $M \in \mathbb{C}[x][\partial]$ s.t.

- ▶ $M = PL$ for some $P \in \mathbb{C}(x)[\partial]$;
- ▶ the origin is an ordinary point of M .

Call M a **desingularized operator** of L .

Desingularization

Given $L \in \mathbb{C}[x][\partial]$, the origin being apparent, find $M \in \mathbb{C}[x][\partial]$ s.t.

- ▶ $M = PL$ for some $P \in \mathbb{C}(x)[\partial]$;
- ▶ the origin is an ordinary point of M .

Call M a **desingularized operator** of L .

A first idea (Fuchs). Assume missing exponents are k_1, \dots, k_ℓ .
Compute the least common left multiple of

$$L, x\partial - k_1, \dots, x\partial - k_\ell$$

in $\mathbb{C}(x)[\partial]$.

Advanced method

Chen, Jaroschek, Kauers and Singer (2013, 2016), construct a desingularized operator M of L s.t.

- ▶ all apparent singularities of L are ordinary points of M ;
- ▶ all singularities of M are non-apparent ones of L ;
- ▶ the degree of leading coeff of M is **minimal**.

Contraction of Ore ideals (Z, 2016)

Theorem. A desingularized operator yields generators of

$$(\mathbb{C}(x)[\partial]L) \cap \mathbb{C}[x][\partial].$$

Contraction of Ore ideals (Z, 2016)

Theorem. A desingularized operator yields generators of

$$(\mathbb{C}(x)[\partial]L) \cap \mathbb{C}[x][\partial].$$

- ▶ Determine the contraction ideals of shift operators
- ▶ The ring of constants can be replaced by a PID

D-finite systems

Notation.

$$\begin{array}{ccc} A_n = \mathbb{C}[x_1, \dots, x_n][\partial_1, \dots, \partial_n] & \subset & R_n = \mathbb{C}(x_1, \dots, x_n)[\partial_1, \dots, \partial_n] \\ \uparrow & & \uparrow \\ \text{Weyl algebra} & & \text{Rational algebra} \end{array}$$

where $\partial_i = \partial/\partial x_i$.

Definition. A left ideal $I \subset R_n$ is **D-finite** if R_n/I is a finite-dimensional vector space over $\mathbb{C}(x_1, \dots, x_n)$.

Assume that G_1, \dots, G_m are generators of I . The system

$$G_i(f) = 0, \quad i = 1, \dots, m.$$

is called a **D-finite system**.

D-finite Gröbner bases

Let \prec_{∂} be a graded term order on $\partial_1^{k_1} \cdots \partial_n^{k_n}$, a finite set $G \subset A_n$ is a Gröbner basis w.r.t. \prec_{∂} .

Definition. G is **D-finite** if $R_n \cdot G$ is D-finite. The set

$$\text{PE}(G) = \left\{ (i_1, \dots, i_n) \mid \partial_1^{i_1} \cdots \partial_n^{i_n} \text{ is not reducible w.r.t. } G \right\}.$$

is called the set of **parametric exponents** of G .


$|\text{PE}(G)|$ is called the **rank** of G .

Example 1

Consider

$$M = \{\partial_1^3, \partial_1^2 \partial_2, \partial_1 \partial_2^2, \partial_2^3\}.$$

Then $PE(M) = \{(0, 0), (1, 0), (0, 1), (2, 0), (1, 1), (0, 2)\}$.



Ordinary points and singularities

Assume that $G \subset A_n$ is a Gröbner basis and its elements are all primitive.

Definition. $c \in \mathbb{C}^n$ is an **ordinary point** of G if c is not a zero of

$$\prod_{g \in G} lc(g).$$

Otherwise, c is a **singularity** of G .

Ordinary points and singularities

Example 1 (cont.) Consider

$$M = \{\partial_1^3, \partial_1^2 \partial_2, \partial_1 \partial_2^2, \partial_2^3\}.$$

where $\prod_{g \in M} \text{lc}(g) = 1$. The origin is an ordinary point of M .

Example 2. Consider

$$G = \{x_2^2 \partial_2 - x_1^2 \partial_1 + x_1 - x_2, \partial_1^2\},$$

where $\prod_{g \in G} \text{lc}(g) = x_2^2$. The origin is a singularity of G .

Formal power series

Let \prec_x be the order induced by \prec_∂ on $x_1^{k_1} \cdots x_n^{k_n}$.

Let $f \in \mathbb{C}[[x_1, \dots, x_n]]$ be of form

$$f = c_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n} + \text{higher terms w.r.t. } \prec_x,$$

where $c_{i_1, \dots, i_n} \in \mathbb{C}$ is nonzero.

Definition. Call (i_1, \dots, i_n) the **initial exponent** of f .

Main result

Let G be a D-finite Gröbner basis and its elements are all primitive.

Theorem. The origin of \mathbb{C}^n is an ordinary point of G



$\forall (i_1, \dots, i_n) \in \text{PE}(G), \exists f \in \mathbb{C}[[x_1, \dots, x_n]]$ with initial exponent (i_1, \dots, i_n) s.t. f is a solution of G .

Main result

Let G be a D-finite Gröbner basis and its elements are all primitive.

Theorem. The origin of \mathbb{C}^n is an ordinary point of G



$\forall (i_1, \dots, i_n) \in \text{PE}(G), \exists f \in \mathbb{C}[[x_1, \dots, x_n]]$ with initial exponent (i_1, \dots, i_n) s.t. f is a solution of G .

Remark. an algorithm for computing formal power series sols of D-finite systems at ordinary points.

Apparent singularities

Assume the origin is a singularity of G .

Definition. The origin is **apparent** if G has $|\text{PE}(G)|$ \mathbb{C} -linearly independent sols in $\mathbb{C}[[x_1, \dots, x_n]]$.

Example 2 (cont.) Consider

$$G = \{x_2^2 \partial_2 - x_1^2 \partial_1 + x_1 - x_2, \partial_1^2\},$$

$\{x_1 + x_2, x_1 x_2\}$ are sols of G . The origin is apparent.

We can decide whether a given point is apparent or not and remove it using “a first idea”.

Example 2 (cont.)

Consider

$$G = \{x_2^2 \partial_2 - x_1^2 \partial_1 + x_1 - x_2, \partial_1^2\},$$

Set

$$S = \{(0, 0), (0, 1), (2, 0), (0, 2)\}.$$

Let $M \subset A_n$ be a Gröbner basis with

$$R_n M = R_n G \cap \left(\bigcap_{(s,t) \in S} R_n \{x_1 \partial_1 - s, x_2 \partial_2 - t\} \right)$$

We find

$$M = \{\partial_1^3, \partial_1^2 \partial_2, \partial_1 \partial_2^2, \partial_2^3\}.$$

The origin is an ordinary point of M .

Conclusion

- ▶ Characterization of ordinary points of D-finite systems
- ▶ Detect and remove apparent singularities of D-finite systems

Conclusion

- ▶ Characterization of ordinary points of D-finite systems
- ▶ Detect and remove apparent singularities of D-finite systems

Remark 1. an algorithm for computing formal power series sols of D-finite systems at apparent singularities.

Conclusion

- ▶ Characterization of ordinary points of D-finite systems
- ▶ Detect and remove apparent singularities of D-finite systems

Remark 1. an algorithm for computing formal power series sols of D-finite systems at apparent singularities.

Remark 2. for arbitrary singularities, Takayama (2003) gives an algorithm by using D-module theory. **No elementary proof!**

Conclusion

- ▶ Characterization of ordinary points of D-finite systems
- ▶ Detect and remove apparent singularities of D-finite systems

Remark 1. an algorithm for computing formal power series sols of D-finite systems at apparent singularities.

Remark 2. for arbitrary singularities, Takayama (2003) gives an algorithm by using D-module theory. **No elementary proof!**

Thanks!